

# UMBILICAL CORD/BELLY BUTTON/NAELSTRING

revision date 18042023

dr hester van der walt


weskus pediatrie / west coast paediatrics

21 voortrekker str, vredenburgh, 7380; tel 0227135149, www.pediater.co.za

pr 0377597, mp 0279102, sama 11741, vat 4530258856, registration 2013/207395/21

---

## Umbilical Cord/Belly Button



Most first-time parents are understandably worried when they notice bleeding or discharge from their new-born's belly button.

Here is a little guide to help you understand when this is normal and when you need to call your doctor.

A brand new recommendation in 2006 is that you no longer need to put alcohol on the umbilical cord. A research study found that it really isn't necessary.

### **BLEEDING**

Most new-borns will have some oozing of blood from the belly button before and especially after the cord falls off. This can either occur right away, or may not appear until a week or more after the cord comes off. It is especially common if the cord comes off within the first week of life from accidental tugging. The blood is usually first noticed during a diaper change when you find some blood on the diaper or T-shirt. This mild oozing of blood can go on for one or two weeks, so don't worry. You may even see blood with every diaper change. You can also pack some gauze over the belly button under the diaper.

**THIS SMALL AMOUNT OF BLEEDING IS NORMAL AND IS**

**NO CAUSE FOR ALARM!**

### **DRAINAGE OR DISCHARGE**

Virtually every belly button will have some yellow or green drainage, both before and after it falls off. It may look like pus, but it is really just mucus. You may notice some on baby's T-shirt or diaper with every diaper change. This may go on for one or two weeks, so don't worry. The drainage will eventually stop. **For excessive oozing, your doctor can help dry up the cord by applying silver nitrate in his/her consulting room. This is rarely necessary.**

## UMBILICAL CORD/BELLY BUTTON/NAELSTRING

revision date 18042023

dr hester van der walt

weskus pediatrie / west coast paediatrics

21 voortrekker str, vredenburgh, 7380; tel 0227135149, www.pediater.co.za

pr 0377597, mp 0279102, sama 11741, vat 4530258856, registration 2013/207395/21

---

### WHEN TO WORRY

If you see **active dripping or pooling of blood that reappears immediately after wiping it away**, then you should pack several pieces of gauze over the belly button. If you do not have gauze, use a baby washcloth or toilet paper. Keep the gauze firmly pressed against the belly button under the diaper, and snap a tight outfit onto baby. Wrap him up snugly in a blanket and wait for 15 minutes. Undress him and carefully check for continued active oozing or dripping. If it has stopped, then keep a close eye on it, and keep gauze packed on it for another day. Carefully check once an hour, even overnight. You will see some blood drying on the gauze. **There is no need to call your doctor.**

**If the active dripping or oozing continues after the 15 minutes, then you should call your doctor right away.**

The only other time you need to worry is when the cord has become **infected**. Here is how to tell:

- ! **The drainage has a very foul odour**
- ! **The skin around the cord has become very red and maybe swollen**
- ! **Baby may or may not have a fever**

Call your doctor if the above symptoms accompany a fever of 38.3C/101F!